

# Fewer than one per cent of Canadians who received a prescription in the past six months say cost is a contributor to non-adherence to prescription medicines

## Prescription Use Survey Summary

Submitted by Nanos to Innovative Medicines Canada & the Canadian Life and Health Insurance Association

Released November 20<sup>th</sup>, 2017



# Key Findings

1. 80% of respondents report having received a prescription in the past six months
2. 93% of respondents who report having received a prescription in the past six months report the prescriptions were taken without exception as prescribed
3. Seven per cent of respondents who report being given a prescription in the last six months did not take it as prescribed. Of those, the top reasons for non-adherence were as follows:
  - 33% say they no longer needed it
  - 23% reported side effects, an allergic reaction or the medication being too strong
  - Seven per cent indicated the medication was too expensive or they couldn't afford it. This shows that less than one per cent of respondents who received a prescription in the last six months did not take the medication(s) as prescribed due to cost
  - Seven per cent forgot to take it
4. Nearly six in ten respondents who report having received a prescription in the last six months and not taking it as prescribed were either not aware (54%) or unsure (three per cent) of public drug programs to help pay for prescription drugs



# The vast majority of respondents who received a prescription in the past six months took it as prescribed

Most respondents report taking prescriptions as prescribed while one in 20 report not taking their prescription medicine according to a new random telephone survey of 4,445 Canadians. Those few who reported personally receiving a prescription but not taking it as prescribed most often said they stopped early/did not fill it/took a smaller dose one time, and frequently said this was due to no longer needing it or side effects.

Readers should note that this research is among Canadian adults who lived in a household where someone had received a prescription medicine in the last six months.



## Prescription use in household

- **More than nine in ten respondents with a child who received a prescription said it was taken as prescribed** – Ninety-four per cent of respondents with a child who received a prescription in the last six months said all prescriptions were taken without exception as prescribed, while five per cent said they were not.
- **Eight in ten respondents reported having received a prescription in the last six months** – Eight in ten respondents (80%) reported they have received a prescription in the past six months, while 20 per cent reported they have not.
- **Majority of respondents who received a prescription in the past six months said it was taken as prescribed** – Over nine in ten (93%) respondents who reported receiving a prescription in the past six months said it was taken as prescribed without exception, while seven per cent said it was not.
- **More than half of the one in 20 respondents who received a prescription and did not take it as prescribed said they did not fill a prescription one time in the last six months** – Fifty-six per cent of respondents who received a prescription in the last six months but did not take it as prescribed said they did not fill a prescription one time, while 40 per cent said they did not fill a prescription more than one time in the past six months. Two per cent were unsure and two per cent refused to answer.
- **Half of the one in 20 respondents who received a prescription and did not take it as prescribed said they stopped taking a prescription early one time in the last six months** – Fifty-two per cent of respondents who received a prescription in the last six months but did not take it as prescribed said they stopped taking the prescription early one time, while 43 per cent said they stopped early more than one time in the past six months. Six per cent were unsure.
- **Four in ten respondents who received a prescription and did not take it as prescribed said they took a smaller dose of a prescription one time in the last six months** – Forty-three per cent of respondents who received a prescription in the last six months but did not take it as prescribed said they took a smaller dose of or took a prescription less often than prescribed one time in the last six months, while 37 per cent said they did this more than once. Nineteen per cent were unsure.



- **Take as needed medications were the most frequently reported medication that respondents did not take as prescribed** – Sixty-four per cent of respondents who received a prescription in the past six months but did not take it as prescribed said the prescription was take as needed medication (ex. pain, antibiotics, digestive issues, anti-inflammatory, etc), while 31 per cent said the prescription was for a serious chronic condition (ex. diabetes, cardiovascular, transplant surgery, osteoporosis, etc), followed by medications to help quality of life (ex. ADHD, RA, etc.)(14%). One per cent said they can't recall/unsure, and seven per cent refused to answer.
- **One in three respondents who received a prescription but did not take it as prescribed cited no longer needing it as the reason** – One third of respondents (33%) who received a prescription in the past six months but did not take it as prescribed said it was because they no longer needed it, followed by side effects (18%), cost (seven per cent), forgot/forgetful (seven per cent), and I take it only when needed (five per cent). Fourteen per cent refused to answer.

#### Prescription medicine expenses

- **Respondents who received a prescription but did not take it as prescribed reported spending an average of \$270 on prescription meds in the last six months** – Asked how much they have spent on average at their own expense on prescription medications, respondents who received a prescription but did not take it as prescribed reported spending an average of \$270.35 in the last six months.
- **Over half of respondents who received a prescription but did not take it as prescribed said they are unaware of any publicly funded programs to help people access prescription medicines** – Just over one in two (54%) respondents who received a prescription medicine but did not take it as prescribed said they are not aware of any publicly funded programs to help people access prescription medicines. Forty-three per cent said they are aware of such programs, while three per cent were unsure.
- **Two in three respondents who received a prescription but did not take it as prescribed say they are covered through a private plan** – Two in three respondents who received a prescription but did not take it as prescribed say they are covered under a work health benefit plan either their own or someone in their family (45%), have health benefits and coverage through a union or professional association (13%), or have health benefits and coverage purchased at their own expense through an insurance carrier (eight per cent). Eighteen per cent said their prescription medicines are paid for by a government public plan, although they are not a government employee, and 12 per cent have no health plan coverage and have to pay for the full price of any prescription needed. Three per cent refused to answer. Please note that each province has a different provincial drug plan, making it difficult to compare findings between provinces.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4,445 Canadians, 18 years of age or older, between May 27<sup>th</sup> and July 10<sup>th</sup>, 2017. Participants were randomly recruited by telephone using live agents and administered a survey. The margin of error for a random survey of 4,445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

The research was commissioned by Innovative Medicines Canada and the Canadian Life and Health Insurance Association (CLHIA)



# Objectives

Innovative Medicines Canada and the Canadian Life and Health Insurance Association (CLHIA) retained Nanos to conduct a survey of Canadians about their use of and behaviours surrounding prescription medicine use.

The objective of the research is to gauge the number of Canadians who are uninsured or underinsured, and whether that has impacted their behaviour in terms of their prescription medicine use. The research also sought to explore the incidence of Canadians who do not take prescription medication(s) as prescribed, and the relationship between that behaviour and the cost of the medication(s).


The survey was conducted among 4,445 Canadians, with two subsamples targeted:

- Canadians with a child(ren) who received a prescription in the last six months (n=590); and,
- Canadians who have personally received a prescription in the last six months (n=2,845).

The following two pages detail the incidence for each of these subgroups within the sample of 4,445 Canadians.


# Canadians - Rollout

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=4445, accurate 1.4 percentage points plus or minus, 19 times out of 20.



# Canadians and children - Rollout

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=4445, accurate 1.4 percentage points plus or minus, 19 times out of 20.



# 1.0 Prescription use in household




# Prescriptions in household

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=4445, accurate 1.4 percentage points plus or minus, 19 times out of 20.

## Net Score

+59.6

Not  
received a  
prescription  
in the past  
six months  
20%



Yes, our  
household  
has received  
a  
prescription  
in the past  
six months  
80%

**\*Note: Charts may not add up to 100 due to rounding**

## Sample = All Canadians

Subgroups	Yes, has received
Atlantic (n=312)	83.7%
Quebec (n=1040)	80.7%
Ontario (n=1720)	79.6%
Prairies (n=302)	81.9%
Alberta (n=491)	81.8%
British Columbia (n=580)	73.9%
Male (n=1925)	76.4%
Female (n=2520)	82.9%
18 to 34 (n=1003)	70.0%
35 to 54 (n=1582)	82.8%
55 plus (n=1860)	84.1%
Under 65 (n=3343)	78.6%
Over 65 (n=1102)	83.8%
\$0-\$29,999 a year (n=658)	80.1%
\$30-\$59,999 a year (n=914)	79.0%
\$60-\$99,999 a year (n=806)	82.5%
\$100K plus a year (n=795)	83.7%

**QUESTION** – In the past six months, has anyone in your household including children, been given a prescription, either new or for a refill, for a medication?


# Prescriptions for children in household

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=4445, accurate 1.4 percentage points plus or minus, 19 times out of 20.

## Net Score

+4.4

No Children  
78%



\*Note: Charts may not add up to 100 due to rounding

**QUESTION – [IF HAS CHILD]** Have any of the children under 18 years of age been given a prescription in the past six months?

## Sample = All Canadians


Subgroups	Yes
Atlantic (n=312)	12.7%
Quebec (n=1040)	13.0%
Ontario (n=1720)	13.3%
Prairies (n=302)	9.8%
Alberta (n=491)	17.8%
British Columbia (n=580)	11.9%
Male (n=1925)	12.0%
Female (n=2520)	14.5%
18 to 34 (n=1003)	24.1%
35 to 54 (n=1582)	18.3%
55 plus (n=1860)	1.1%
Under 65 (n=3343)	17.1%
Over 65 (n=1102)	0.3%
\$0-\$29,999 a year (n=658)	6.9%
\$30-\$59,999 a year (n=914)	9.1%
\$60-\$99,999 a year (n=806)	15.7%
\$100K plus a year (n=795)	23.8%

# Prescriptions for children taken as prescribed

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=555, accurate 4.2 percentage points plus or minus, 19 times out of 20.

**Net Score**

**+89.1**



**\*Note: Charts may not add up to 100 due to rounding**

**QUESTION – [IF CHILD PRESCRIPTION IN PAST 6 MONTHS]** All prescriptions for children under 18 years of age were taken without exception as prescribed?

Please note that questions were asked following this question related to children not having taken medications as prescribed but the responding sample size of each was less than 30 and they have not been reported on. See the methodology for the questions and see the Appendix for the full questionnaire.

**Sample = Canadians with a child who has received a prescription in past 6 months**


Subgroups	Yes
Atlantic (n=35)	100.0%
Quebec (n=125)	93.2%
Ontario (n=221)	93.9%
Prairies (n=27)*	
Alberta (n=82)	91.6%
British Columbia (n=65)	96.1%
Male (n=215)	94.3%
Female (n=340)	94.1%
18 to 34 (n=245)	93.0%
35 to 54 (n=290)	95.8%
55 plus (n=20)*	
Under 65 (n=552)	94.3%
Over 65 (n=3)*	
\$0-\$29,999 a year (n=40)	95.4%
\$30-\$59,999 a year (n=76)	94.5%
\$60-\$99,999 a year (n=124)	95.7%
\$100K plus a year (n=183)	93.6%

**\*Masked due to sample size being 30 or less**

# Personal prescription use

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=3590, accurate 1.6 percentage points plus or minus, 19 times out of 20.

**Net Score**  
**+60.6**



**\*Note: Charts may not add up to 100 due to rounding**

**QUESTION – [IF SOMEONE IN HOUSEHOLD RECEIVED A PRESCRIPTION IN THE PAST SIX MONTHS] Have you been given a prescription in the past six months?**

**Sample = Those with someone in household who received prescription in past 6 months**


Subgroups	Yes
Atlantic (n=264)	84.7%
Quebec (n=845)	81.9%
Ontario (n=1388)	78.3%
Prairies (n=251)	85.4%
Alberta (n=409)	80.9%
British Columbia (n=433)	76.7%
Male (n=1485)	78.8%
Female (n=2105)	81.5%
18 to 34 (n=707)	71.2%
35 to 54 (n=1318)	75.4%
55 plus (n=1565)	89.7%
Under 65 (n=2665)	76.6%
Over 65 (n=925)	91.6%
\$0-\$29,999 a year (n=532)	88.3%
\$30-\$59,999 a year (n=731)	80.9%
\$60-\$99,999 a year (n=673)	80.0%
\$100K plus a year (n=673)	74.0%

# Personal prescriptions taken as prescribed

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=2904, accurate 1.8 percentage points plus or minus, 19 times out of 20.

**Net Score**

+85.6



**\*Note: Charts may not add up to 100 due to rounding**


**QUESTION – [IF PERSONAL PRESCRIPTION IN PAST 6 MONTHS]** You personally took all prescriptions without exception as prescribed?

**Sample = Canadians who received a prescription in the past 6 months**

Subgroups	Yes
Atlantic (n=223)	95.7%
Quebec (n=699)	87.7%
Ontario (n=1096)	94.7%
Prairies (n=214)	93.1%
Alberta (n=335)	95.0%
British Columbia (n=337)	92.6%
Male (n=1181)	92.6%
Female (n=1723)	92.9%
18 to 34 (n=503)	88.2%
35 to 54 (n=998)	93.0%
55 plus (n=1403)	94.7%
Under 65 (n=2057)	91.8%
Over 65 (n=847)	95.2%
\$0-\$29,999 a year (n=471)	93.6%
\$30-\$59,999 a year (n=595)	91.9%
\$60-\$99,999 a year (n=543)	92.2%
\$100K plus a year (n=504)	92.9%

# Personal prescriptions stopped early

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=153, accurate 8.0 percentage points plus or minus, 19 times out of 20.



**QUESTION – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED]** You personally stopped taking the prescription early \_\_\_\_ occasions in the past six months.


**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

Subgroups	One time
Atlantic (n=7)*	
Quebec (n=65)	51.4%
Ontario (n=41)	53.3%
Prairies (n=13)*	
Alberta (n=11)*	
British Columbia (n=16)*	
Male (n=70)	51.6%
Female (n=83)	51.4%
18 to 34 (n=44)	50.3%
35 to 54 (n=52)	57.1%
55 plus (n=57)	47.5%
Under 65 (n=121)	53.2%
Over 65 (n=32)	44.2%
\$0-\$29,999 a year (n=22)*	
\$30-\$59,999 a year (n=34)	41.7%
\$60-\$99,999 a year (n=34)	52.3%
\$100K plus a year (n=24)*	

**\*Masked due to sample size being 30 or less**

# Personal prescriptions not filled

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=111, accurate 9.4 percentage points plus or minus, 19 times out of 20.



\*Note: Charts may not add up to 100 due to rounding

**QUESTION – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED] You personally received but did not to fill a prescription \_\_\_\_\_ occasions in the past 6 months**


**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

Subgroups	One time
Atlantic (n=7)*	
Quebec (n=63)	48.7%
Ontario (n=21)*	
Prairies (n=7)*	
Alberta (n=11)*	
British Columbia (n=8)*	
Male (n=43)	49.6%
Female (n=68)	59.8%
18 to 34 (n=30)*	
35 to 54 (n=42)	52.9%
55 plus (n=39)	61.0%
Under 65 (n=85)	56.3%
Over 65 (n=26)*	
\$0-\$29,999 a year (n=14)*	
\$30-\$59,999 a year (n=27)*	
\$60-\$99,999 a year (n=27)*	
\$100K plus a year (n=18)*	

\*Masked due to sample size being 30 or less

# Smaller doses for personal prescription

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=145, accurate 8.3 percentage points plus or minus, 19 times out of 20.



**QUESTION – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED]** You personally took a smaller dose or less often than prescribed \_\_\_\_ occasions in the past 6 months

**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

Subgroups	One time
Atlantic (n=7)*	
Quebec (n=70)	53.0%
Ontario (n=35)	32.9%
Prairies (n=7)*	
Alberta (n=8)*	
British Columbia (n=18)*	
Male (n=60)	45.2%
Female (n=85)	41.9%
18 to 34 (n=40)	52.2%
35 to 54 (n=49)	48.1%
55 plus (n=56)	30.6%
Under 65 (n=114)	44.3%
Over 65 (n=31)	39.7%
\$0-\$29,999 a year (n=21)*	
\$30-\$59,999 a year (n=38)	45.4%
\$60-\$99,999 a year (n=27)*	
\$100K plus a year (n=23)*	

**\*Masked due to sample size being 30 or less**

# Personal prescriptions not taken as prescribed

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=2904, accurate 1.8 percentage points plus or minus, 19 times out of 20.

	No
You personally took all prescriptions without exception as prescribed	n=206
	At least 1 occasion in the last six months
You personally stopped taking the prescription early in the past six months.	n=148
You personally received but did not to fill a prescription in the past 6 months	n=107
You personally took a smaller dose or less often than prescribed in the past 6 months	n=118



**QUESTION – [IF PERSONAL PRESCRIPTION IN PAST 6 MONTHS]** You personally took all prescriptions without exception as prescribed?

# Personal prescriptions not taken as prescribed

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=174, accurate 1.8 percentage points plus or minus, 19 times out of 20.

	Frequency (n=174)
Has not taken as prescribed in <b><u>ONE</u></b> of the following ways: took a smaller dose OR did not fill OR stopped taking early	30.3%
Has not taken as prescribed in <b><u>TWO</u></b> of the following ways: took a smaller dose OR did not fill OR stopped taking early	30.4%
Has not taken as prescribed in all <b><u>THREE</u></b> of the following ways: took a smaller dose OR did not fill OR stopped taking early	39.4%

\*Based on multiple mentions

**QUESTION** – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED] You personally stopped taking the prescription early \_\_\_\_ occasions in the past six months.

**QUESTION** – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED] You personally received but did not to fill a prescription \_\_\_\_ occasions in the past 6 months

**QUESTION** – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED] You personally took a smaller dose or less often than prescribed \_\_\_\_ occasions in the past 6 months

# Reasons for personal prescriptions

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=178, accurate 7.4 percentage points plus or minus, 19 times out of 20.

Top responses	Frequency (n=178)*
Pain medication	20.6%
Antibiotics	13.2%
Digestive issues	10.3%
Depression	9.3%
Anti-inflammatory	9.1%
Anxiety	5.3%
Allergies	4.9%
Cholesterol	4.7%
Sleep aids	4.6%
Medication for diabetes	4.5%
Blood pressure	4.4%
Refuse	7.0%
Unsure	0.6%

**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

\*Based on multiple mentions

**QUESTION –** [IF PERSONAL PRESCRIPTION IN PAST 6 MONTHS AND NOT TAKEN AS PRESCRIBED] In general, what were those prescriptions for? [Open-ended]

# Reasons for personal prescriptions - grouped

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=178, accurate 7.4 percentage points plus or minus, 19 times out of 20.

Top responses	Frequency (n=178)*
Take as needed medications	64.1%
Medications for serious chronic conditions	30.5%
Medications to help quality of life	13.9%
Can't recall/unsure	1.2%
Refuse	7.1%
Other	5.0%

**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

\*Based on multiple mentions

**QUESTION – [IF PERSONAL PRESCRIPTION IN PAST 6 MONTHS AND NOT TAKEN AS PRESCRIBED]** In general, what were those prescriptions for? [Open-ended]

# Reasons for not taking personal prescription as prescribed

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=191, accurate 7.2 percentage points plus or minus, 19 times out of 20.

	Frequency (n=191)*
No longer needed it	33.0%
Side effects	18.2%
Too expensive/couldn't afford it	7.2%
Forgot/forgetful	7.0%
I take it only when needed	5.2%
Allergic reaction	2.9%
Medication was changed	2.9%
Did not help my issues	2.5%
Meds were too strong	2.3%
I did not want to take it	2.0%
I prefer natural medicines	1.5%
No reason	0.9%
Other	0.5%
Unsure	0.5%
Refuse	13.6%

\*Based on multiple mentions

**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

Please note that following this question, a question regarding the cost of prescriptions was asked of those who indicated prescriptions were not taken as prescribed due to cost, but the responding sample size was less than 30 and they have not been reported on. See the methodology for the question and see the Appendix for the full questionnaire.

**QUESTION** – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED] What was the one reason for [not filling/stopping early/taking a smaller dose of] your personal prescription on any occasion. [Open-ended]

# Reasons for not taking personal prescription as prescribed - Grouped

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=191, accurate 7.2 percentage points plus or minus, 19 times out of 20.

	Frequency (n=191)*
No longer needed it	33.0%
Side effects/Allergic reaction/ Meds were too strong	23.4%
Too expensive/couldn't afford it	7.2%
Forgot/forgetful	7.0%
I take it only when needed	5.2%
Medication was changed	2.9%
Did not help my issues	2.5%
I did not want to take it	2.0%
I prefer natural medicines	1.5%
No reason	0.9%
Other	0.5%
Unsure	0.5%
Refuse	13.6%

**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

Please note that following this question, a question regarding the cost of prescriptions was asked of those who indicated prescriptions were not taken as prescribed due to cost, but the responding sample size was less than 30 and they have not been reported on. See the methodology for the question and see the Appendix for the full questionnaire.

**\*Based on multiple mentions**

**QUESTION – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED]** What was the one reason for [not filling/stopping early/taking a smaller dose of] your personal prescription on any occasion. [Open-ended]

## 2.0 Prescription medicine expenses



# Personal expenses on prescriptions

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=147 accurate 8.2 percentage points plus or minus, 19 times out of 20.

Mean	Median
\$270.35	\$100.00

Ten most frequent responses	Frequency (n=147)
\$100	13.4%
\$200	10.9%
\$50	9.6%
\$30	9.5%
\$300	6.6%
\$20	6.5%
\$150	3.9%
\$600	2.3%
\$80	2.2%
\$500	2.2%

**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**


Subgroups	Mean
Under 65 (n=150)	\$281.86
Over 65 (n=41)	\$224.49
\$0-\$29,999 a year (n=30)*	
\$30-\$59,999 a year (n=47)	\$205.19
\$60-\$99,999 a year (n=41)	\$306.64
\$100K plus a year (n=34)	\$330.96

**\*Masked due to sample size being 30 or less**

**QUESTION – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED]** In the past six months, about how much was spent in total, at your own expense, on prescription medicines? [Open-ended]

# Awareness of publicly funded prescription programs

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=203, accurate 7.0 percentage points plus or minus, 19 times out of 20.



**\*Note: Charts may not add up to 100 due to rounding**

**QUESTION – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED] Are you aware or not aware of any publicly funded programs to help people access prescription medicines?**

**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

Subgroups	Aware
Atlantic (n=10)*	
Quebec (n=83)	37.4%
Ontario (n=55)	38.3%
Prairies (n=15)*	
Alberta (n=15)*	
British Columbia (n=25)*	
Male (n=85)	44.3%
Female (n=118)	42.2%
18 to 34 (n=59)	40.6%
35 to 54 (n=68)	41.9%
55 plus (n=76)	47.2%
Under 65 (n=161)	42.7%
Over 65 (n=42)	45.5%
\$0-\$29,999 a year (n=30)*	
\$30-\$59,999 a year (n=47)	32.5%
\$60-\$99,999 a year (n=41)	56.6%
\$100K plus a year (n=34)	46.9%

**\*Masked due to sample size being 30 or less**

# Health benefit plan coverage

Source: Nanos Research, RDD dual frame telephone random survey, May 27<sup>th</sup> to July 10<sup>th</sup>, 2017, n=203, accurate 7.0 percentage points plus or minus, 19 times out of 20.

	Frequency (n=203)
I am covered through a work health benefit plan which is either my own or someone in my family	44.6%
My prescription medicines are paid for by a government public plan but I am not a government employee	18.4%
I have health plan benefits and coverage through a union or professional association	12.8%
<u>I have no health plan coverage</u> and all the time have to pay for the full price of any prescription I need	11.8%
I have health plan benefits and coverage which has been purchased, at our own expense through an insurance carrier	8.1%
Other	0.8%
Refuse	3.4%
	Frequency (n=203)
I am covered through a work health benefit plan <b>OR</b> I have health plan benefits and coverage through a union or professional association <b>OR</b> I have health plan benefits and coverage which has been purchased, at our own expense through an insurance carrier	65.5%

**\*Note: Charts may not add up to 100 due to rounding**


**QUESTION** – [IF PERSONAL PRESCRIPTION NOT TAKEN AS PRESCRIBED] I am going to read a series of statements about your health benefit plan coverage and I would like you to tell me which best describes you [RANDOMIZE]

Please note that following this question, a question regarding reasons for not having insurance or government benefits was asked, but the responding sample size of each was less than 30 and they have not been reported on. See the methodology for the question and see the Appendix for the full questionnaire.

**Sample = Canadians who received a prescription in the past six months and did not take it as prescribed**

Subgroups	Work health benefit plan
Atlantic (n=10)*	
Quebec (n=83)	46.2%
Ontario (n=55)	39.6%
Prairies (n=15)*	
Alberta (n=15)*	
British Columbia (n=25)*	
Male (n=85)	45.8%
Female (n=118)	43.6%
18 to 34 (n=59)	51.0%
35 to 54 (n=68)	63.0%
55 plus (n=76)	20.6%
Under 65 (n=161)	52.6%
Over 65 (n=42)	8.7%
\$0-\$29,999 a year (n=30)*	
\$30-\$59,999 a year (n=47)	37.6%
\$60-\$99,999 a year (n=41)	60.9%
\$100K plus a year (n=34)	63.4%

**\*Masked due to sample size being 30 or less**



# Methodology

# Methodology

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4,445 Canadians, 18 years of age or older, between May 27<sup>th</sup> and July 10<sup>th</sup>, 2017. Participants were randomly recruited by telephone using live agents and administered a survey. The results were statistically checked and weighted by age and gender using the latest Census information and the sample is geographically stratified to be representative of Canada.

Individuals were randomly called using random digit dialing with a maximum of five call backs.

The margin of error for a random survey of 4,445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

The research was commissioned by Innovative Medicines Canada and the Canadian Life and Health Insurance Association (CLHIA)

Note: Charts may not add up to 100 due to rounding.

# Methodology

**The following questions received fewer than 30 responses:**

**Question 5** - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age received a prescription but the prescription was not filled (occasions in the past 6 months) [OPEN]

**Question 6** - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age stopped take a prescriptions early (occasions in the past 6 months) [OPEN]

**Question 7** - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age took a smaller dose of the medication or less than prescribed (occasions in the past 6 months) [OPEN]

**Question 8** – [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND CHILD’S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In general, what were those prescriptions for? [OPEN]

**Question 11** – [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND CHILD’S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] What was the one reason for [not filling/stopping early/taking a smaller dose of] the prescription of the child under 18 years of age on any occasion? [OPEN]

**Question 12** - [IF COST IS THE REASON A CHILD’S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN]

**Question 22** - [IF COST IS THE REASON FOR PRESCRIPTION FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)? [OPEN]

**Question 26** - [IF PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN AND NO INSURANCE FOR PRESCRIPTION MEDICINES] Why do you not have any insurance or government benefits for prescription medication? [OPEN]

# Technical Note

Element	Description	Element	Description
Organization who commissioned the research	Innovative Medicines Canada and CLHIA	Weighting of Data	The results were weighted by age and gender using the latest Census information (2014) and the sample is geographically stratified to ensure a distribution across all regions of Canada. See tables for full weighting disclosure
Final Sample Size	4,445 Randomly selected individuals.	Screening	Screening ensured potential respondents did not work in the market research industry or in the advertising industry prior to administering the survey to ensure the integrity of the data.
Margin of Error	±1.4 percentage points, 19 times out of 20.	Excluded Demographics	Individuals younger than 18 years old; individuals without land or cell lines could not participate.
Mode of Survey	RDD dual frame (land- and cell-lines) telephone survey	Stratification	By age and gender using the latest Census information (2014) and the sample is geographically stratified to be representative of Canada. Smaller areas such as Atlantic Canada were marginally oversampled to allow for a minimum regional sample.
Sampling Method Base	The sample included both land- and cell-lines RDD (Random Digit Dialed) across Canada.	Estimated Response Rate	Eight percent, consistent with industry norms.
Demographics (Captured)	Atlantic Canada, Quebec, Ontario, Prairies, British Columbia; Men and Women; 18 years and older. Six digit postal code was used to validate geography.	Question Order	Question order in the preceding report reflects the order in which they appeared in the original questionnaire.
Fieldwork/Validation	Live interviews with live supervision to validate work as per the MRIA Code of Conduct	Question Content	Data with more than 30 respondents in the sample are in the report. Please see the methodology for a complete list of the questions excluded from the report due to small sample sizes, and see the Appendix B for the full questionnaire.
Number of Calls	Maximum of five call backs.	Question Wording	The questions in the preceding report are written exactly as they were asked to individuals.
Time of Calls	Individuals were called between 12-5:30 pm and 6:30-9:30pm local time for the respondent.	Survey Company	Nanos Research
Field Dates	May 27 <sup>th</sup> to July 10 <sup>th</sup> , 2017.	Contact	Contact Nanos Research for more information or with any concerns or questions. <a href="http://www.nanosresearch.com">http://www.nanosresearch.com</a> Telephone:(613) 234-4666 ext. Email: info@nanosresearch.com.
Language of Survey	The survey was conducted in both English and French.		

# About Nanos

Nanos is one of North America's most trusted research and strategy organizations. Our team of professionals is regularly called upon by senior executives to deliver superior intelligence and market advantage whether it be helping to chart a path forward, managing a reputation or brand risk or understanding the trends that drive success. Services range from traditional telephone surveys, through to elite in-depth interviews, online research and focus groups. Nanos clients range from Fortune 500 companies through to leading advocacy groups interested in understanding and shaping the public landscape. Whether it is understanding your brand or reputation, customer needs and satisfaction, engaging employees or testing new ads or products, Nanos provides insight you can trust.



View our brochure

## **Nanos Research**

North America Toll-free

1.888.737.5505

[info@nanosresearch.com](mailto:info@nanosresearch.com)



# Tabulations



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 1 - In the past six months, has anyone in your household, including children, been given a prescription, either new or for a refill for a medication?	Total	Unwgt N	4445	312	1040	1720	302	491	580	1925	2520	1003	1582	1860
		Wgt N	4444	312	1041	1719	302	491	580	2148	2296	1212	1515	1717
	Yes	%	79.8	83.7	80.7	79.6	81.9	81.8	73.9	76.4	82.9	70.0	82.8	84.1
	No	%	20.2	16.3	19.3	20.4	18.1	18.2	26.1	23.6	17.1	30.0	17.2	15.9

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 2 - How many children under 18 years of age are in your household? [OPEN]	Total	Unwgt N	4445	312	1040	1720	302	491	580	1925	2520	1003	1582	1860
		Wgt N	4444	312	1041	1719	302	491	580	2148	2296	1212	1515	1717
	No/zero children	%	76.3	79.5	76.7	76.1	80.0	67.9	79.7	78.0	74.7	59.5	67.6	95.9
	One child	%	8.8	8.4	8.6	9.2	5.7	11.5	7.5	8.5	9.1	12.2	14.2	1.6
	Two children	%	9.2	7.5	9.8	8.9	7.2	12.3	8.8	8.5	10.0	17.7	11.9	.9
	Three children	%	3.3	2.6	3.1	3.3	3.7	5.0	2.2	2.9	3.6	6.5	3.9	.4
	Four children	%	1.2	1.5	1.4	1.0	.6	2.2	.6	1.1	1.3	2.6	1.3	.1
	Refuse	%	1.2	.6	.5	1.4	2.7	1.2	1.2	1.0	1.3	1.4	1.1	1.1

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



## 2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 3 - [IF ONE OR MORE CHILDREN] Have any of the children under 18 years of age been given a prescription in the past six months?	Total	Unwgt N	4445	312	1040	1720	302	491	580	1925	2520	1003	1582	1860
		Wgt N	4444	312	1041	1719	302	491	580	2148	2296	1212	1515	1717
	0 children	%	77.5	80.1	77.2	77.5	82.8	69.1	80.9	79.1	76.0	60.9	68.6	97.0
	Yes	%	13.3	12.7	13.0	13.3	9.8	17.8	11.9	12.0	14.5	24.1	18.3	1.1
	No	%	8.9	6.4	9.8	8.8	7.5	12.8	6.8	8.7	9.2	14.5	12.5	1.8
	Unsure	%	.3	.8	.0	.4	.0	.4	.4	.3	.3	.4	.5	.0

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 4 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS] All prescriptions for children under 18 years of age were taken without exception as prescribed?	Total	Unwgt N	555	35	125	221	27	82	65	215	340	245	290	20
		Wgt N	588	40	134	228	29	87	69	257	331	292	277	19
	Yes	%	94.2	100.0	93.2	93.9		91.6	96.1	94.3	94.1	93.0	95.8	
	No	%	5.1	.0	6.8	5.0		8.4	1.3	5.2	5.0	6.1	3.9	
	Unsure	%	.7	.0	.0	1.1		.0	2.5	.5	.9	.9	.3	

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 5 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age received a prescription but the prescription was not filled (occasions in the past 6 months) [OPEN]	Total	Unwgt N	15	0	7	6	0	1	1	7	8	9	6	0
		Wgt N	17	0	8	6	0	1	1	9	8	11	6	0
	1	%												
	2	%												
	4	%												
	Unsure	%												
	No answer	%												

\*Masked due to small sample size.

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 6 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age stopped take a prescriptions early (occasions in the past 6 months) [OPEN]	Total	Unwgt N	21	0	6	7	1	7	0	8	13	12	8	1
		Wgt N	23	0	7	7	1	7	0	10	13	14	7	1
	1	%												
	2	%												
	Unsure	%												

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 7 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age took a smaller dose of the medication or less than prescribed (occasions in the past 6 months) [OPEN]	Total	Unwgt N	14	0	7	3	1	3	0	7	7	8	6	0
		Wgt N	15	0	8	3	1	3	0	9	7	10	6	0
	1	%												
	2	%												
	10	%												
	30	%												
	Unsure	%												

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 8 – [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In general, what were those prescriptions for? [OPEN]	Total	Unwgt N	22	0	8	7	1	5	1	8	14	12	10	0
		Wgt N	23	0	9	7	1	5	1	10	14	14	9	0
	Vitamin deficiency	%												
	Antibiotics	%												
	Asthma	%												
	ADHD	%												
	Acne medication	%												
	Migraines	%												
	Ear infection	%												
	Skin condition/rash	%												
	Dyslexia	%												
	Mosquito bite	%												
	Chronic disease	%												
	Buccal infection	%												
	Lung infection	%												
	As a precaution	%												
	Anti-depressants	%												
	Tonsils	%												

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

		Region							Gender		Age		
		Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Cold relief	%												
Allergies	%												
Refuse	%												

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



## 2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

			Region							Gender		Age		
			Canada											
			2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 11 – [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] What was the one reason for [not filling/stopping early/taking a smaller dose of] the prescription of the child under 18 years of age on any occasion? [OPEN]	Total	Unwgt N	28	0	8	11	1	7	1	11	17	15	12	1
		Wgt N	30	0	9	11	1	7	1	13	16	18	11	1
	Forgot/forgetful	%												
	We already had some reason for [not filling/stopping	%												
	Wrong diagnosis	%												
	Their condition improved/no longer needed	%												
	To see how they would respond	%												
	Child chose not to take it	%												
	Doctor recommendation	%												
	It was not working	%												
	Allergic reaction to medication	%												
	It was unnecessary	%												
	Unsure	%												
	Refuse	%												

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



## 2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

**Question 12 – [IF COST IS THE REASON A CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN] \* Region**

Mean

Question 12 - [IF COST IS THE REASON A CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN]	
Region	
Quebec	
Total	

\*Masked due to small sample size.

**Question 12 - [IF COST IS THE REASON A CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN] \* Gender**

Mean

Question 12 - [IF COST IS THE REASON A CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN]	
Gender	
Female	
Total	

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 13 – [IF SOMEONE IN HOUSEHOLD HAS BEEN GIVEN PRESCRIPTION IN LAST SIX MONTHS] Have you been given a prescription in the past six months?	Total	Unwgt N	3590	264	845	1388	251	409	433	1485	2105	707	1318	1565
		Wgt N	3547	261	840	1368	247	402	428	1642	1904	849	1254	1444
	Yes	%	80.2	84.7	81.9	78.3	85.4	80.9	76.7	78.8	81.5	71.2	75.4	89.7
	No	%	19.6	15.3	18.1	21.4	14.6	18.8	23.3	21.1	18.4	28.5	24.6	10.1
	Unsure	%	.1	.0	.0	.2	.0	.0	.0	.1	.1	.0	.0	.2
	Refuse	%	.1	.0	.0	.1	.0	.3	.0	.1	.1	.3	.0	.0

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 14 – [IF SOMEONE IN HOUSEHOLD HAS BEEN GIVEN PRESCRIPTION IN LAST SIX MONTHS AND PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS] You personally took all prescriptions without exception as prescribed	Total	Unwgt N	2904	223	699	1096	214	335	337	1181	1723	503	998	1403
		Wgt N	2845	221	688	1071	211	325	329	1294	1551	604	946	1295
	Yes	%	92.8	95.7	87.7	94.7	93.1	95.0	92.6	92.6	92.9	88.2	93.0	94.7
	No	%	7.2	4.3	12.3	5.3	6.9	5.0	7.4	7.4	7.1	11.8	7.0	5.3

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 15 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally stopped taking the prescription early (occasions in the past 6 months) [OPEN]	Total	Unwgt N	153	7	65	41	13	11	16	70	83	44	52	57
		Wgt N	157	7	65	43	13	12	16	79	78	53	50	53
	1	%	51.5		51.4	53.3				51.6	51.4	50.3	57.1	47.5
	2	%	28.9		37.5	17.2				24.2	33.7	26.8	25.1	34.7
	3	%	5.2		4.6	9.5				6.8	3.5	1.9	4.3	9.3
	4	%	.8		.0	3.0				1.7	.0	2.5	.0	.0
	5	%	.8		1.8	.0				.0	1.5	2.2	.0	.0
	6	%	.8		.0	3.0				1.7	.0	2.5	.0	.0
	8	%	.6		.0	.0				1.1	.0	.0	.0	1.7
	10	%	1.5		.0	3.0				2.9	.0	2.5	.0	1.8
	12	%	.6		.0	.0				.0	1.2	.0	1.8	.0
	15	%	.8		1.9	.0				1.5	.0	2.3	.0	.0
	20	%	2.1		.0	2.7				.0	4.2	4.4	1.8	.0
	30	%	.8		.0	.0				1.7	.0	.0	2.6	.0
	Unsure	%	5.7		2.8	8.0				6.9	4.4	4.9	7.3	4.9

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 16 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally received but did not fill a prescription (occasions in the past 6 months) [OPEN]	Total	Unwgt N	111	1	63	21	7	11	8	43	68	30	42	39
		Wgt N	112	1	63	22	6	13	8	48	64	36	41	35
	1	%	55.5		48.7					49.6	59.8		52.9	61.0
	2	%	34.2		50.0					34.2	34.2		39.7	35.9
	3	%	5.0		.0					8.0	2.8		2.0	3.1
	4	%	1.2		.0					2.8	.0		3.3	.0
	Unsure	%	1.9		1.4					2.8	1.3		2.1	.0
	Refuse	%	2.2		.0					2.8	1.8		.0	.0

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 17 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally took a smaller dose or less often than prescribed (occasions in the past 6 months) [OPEN]	Total	Unwgt N	145	7	70	35	7	8	18	60	85	40	49	56
		Wgt N	146	7	71	36	7	8	17	67	79	48	47	51
	1	%	43.4		53.0	32.9				45.2	41.9	52.2	48.1	30.6
	2	%	19.5		28.9	.0				14.1	24.0	17.2	19.0	22.0
	3	%	2.0		1.2	3.0				3.0	1.1	.0	.0	5.8
	5	%	1.1		.0	2.3				.0	2.1	.0	.0	3.3
	6	%	1.2		.0	2.2				1.4	1.0	.0	1.7	1.9
	7	%	.8		.0	3.0				1.6	.0	.0	.0	2.2
	10	%	6.4		6.4	5.9				10.1	3.2	5.2	11.0	3.2
	12	%	.6		.0	.0				.0	1.0	.0	.0	1.6
	15	%	.8		.0	3.0				1.6	.0	.0	.0	2.2
	20	%	2.3		.0	6.9				1.9	2.6	5.2	.0	1.6
	21	%	.6		.0	2.3				.0	1.1	.0	.0	1.7
	30	%	2.1		.0	2.2				2.0	2.3	2.1	4.5	.0
	Unsure	%	18.5		8.8	36.0				17.1	19.6	15.6	15.6	23.9
	Refuse	%	.8		1.7	.0				1.8	.0	2.5	.0	.0

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

		Region							Gender		Age		
		Canada 2017-06	Atlantic	Quebec	Ontario	Prairies	Alberta	British Columbia	Male	Female	18 to 34	35 to 54	55 plus
Total	Unwgt N	174	8	76	42	14	14	20	76	98	52	60	62
	Wgt N	178	8	77	44	14	16	20	85	93	62	59	57
One response	%	30.3		21.0	34.0				33.2	27.5	36.1	30.3	23.8
Two responses	%	30.4		13.3	47.6				30.9	29.9	31.7	24.0	35.4
Three responses	%	39.4		65.6	18.4				35.9	42.6	32.1	45.7	40.8

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 18 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In general, what were those personal prescriptions for? [OPEN]	Total	Unwgt N	178	10	76	46	14	10	22	71	107	50	59	69
		Wgt N	180	9	77	47	14	10	22	79	100	60	57	63
	Antibiotics	%	13.2		11.9	18.1				13.3	13.1	22.5	4.6	12.2
	Digestive issues	%	10.3		2.7	22.9				11.3	9.5	4.2	13.9	13.0
	Pain medication	%	20.6		24.9	5.8				21.6	19.8	11.5	24.4	25.8
	Infection	%	.7		.0	2.8				1.7	.0	2.2	.0	.0
	Anxiety	%	5.3		5.7	.0				4.5	5.9	8.1	6.7	1.5
	Incontinence	%	.9		1.1	1.8				.0	1.7	.0	1.5	1.3
	Anti-inflammatory	%	9.1		13.2	6.8				10.2	8.4	11.9	8.8	6.9
	Depression	%	9.3		10.2	6.8				4.0	13.4	11.0	10.0	7.0
	Hypertension	%	3.3		7.7	.0				3.8	2.9	2.0	.0	7.6
	Allergies	%	4.9		4.6	7.0				5.7	4.3	10.1	3.5	1.3
	Fibromyalgia	%	.5		1.1	.0				.0	.9	.0	1.5	.0
	Can't recall	%	.6		.0	2.2				1.3	.0	.0	1.9	.0
	ADHD	%	1.4		1.6	2.8				3.2	.0	4.2	.0	.0
	Cholesterol	%	4.7		6.0	1.8				4.3	5.0	2.4	3.0	8.4
	Skin condition	%	2.6		4.9	.0				3.7	1.8	.0	.0	7.5
	Osteoporosis	%	1.0		1.2	.0				.0	1.7	.0	.0	2.7

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

		Region							Gender		Age		
		Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Headache	%	.5		1.1	.0				.0	.9	.0	1.5	.0
Athlete's Foot	%	.5		.0	.0				1.2	.0	.0	.0	1.5
Muscle spasms	%	2.6		.0	5.1				4.7	1.0	3.9	2.3	1.7
Heart condition	%	.5		.0	.0				.0	.8	.0	.0	1.3
Hemorrhoids	%	.5		.0	.0				1.2	.0	.0	.0	1.5
Rheumatoid arthritis	%	1.3		.0	1.8				.0	2.4	.0	.0	3.8
Abscessed tooth	%	.5		.0	.0				.0	.9	.0	1.6	.0
Flu/cold	%	2.4		1.6	6.5				2.9	2.0	4.0	3.3	.0
Eye drops	%	.5		.0	.0				.0	.9	.0	.0	1.5
Puffer for asthma	%	2.2		.0	4.5				2.8	1.8	2.2	.0	4.2
Hormones	%	1.6		3.8	.0				.0	2.9	2.0	1.5	1.4
Sleep aids	%	4.6		6.8	4.5				5.6	3.8	4.0	8.8	1.4
Blood thinner	%	2.4		1.3	4.0				4.4	.8	.0	1.4	5.5
Alzheimer's	%	.5		.0	.0				1.2	.0	.0	.0	1.5
Blood pressure	%	4.4		4.3	7.5				5.6	3.4	4.1	3.3	5.6
Medication for lung condition	%	.5		.0	.0				1.1	.0	.0	.0	1.4
Medication for diabetes	%	4.5		2.6	9.2				5.6	3.7	4.2	6.9	2.7
Oral contraceptives	%	.7		1.5	.0				.0	1.2	2.0	.0	.0
Hyperthyroidism	%	2.0		2.3	.0				.0	3.5	.0	1.7	4.1

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

		Region							Gender		Age		
		Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Medication for after transplant surgery	%	1.0		2.2	.0				.0	1.7	.0	3.0	.0
Iron supplements	%	1.0		.0	.0				.0	1.7	.0	.0	2.8
Central nervous system	%	.5		1.2	.0				.0	.9	.0	.0	1.4
Medicated shampoo	%	.6		.0	.0				.0	1.0	1.7	.0	.0
Fetal malformation	%	.7		1.5	.0				.0	1.2	2.0	.0	.0
Other	%	5.6		6.6	8.8				4.3	6.6	6.0	1.9	8.5
Unsure	%	.6		.0	.0				.0	1.2	1.9	.0	.0
Refuse	%	7.6		6.5	6.9				9.2	6.3	6.5	8.9	7.3

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-06	Atlantic	Quebec	Ontario	Prairies	Alberta	British Columbia	Male	Female	18 to 34	35 to 54	55 plus
Question 18 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In general, what were those personal prescriptions for? [OPEN]	Total	Unwgt N	178	10	76	46	14	10	22	71	107	50	59	69
		Wgt N	180	9	77	47	14	10	22	79	100	60	57	63
	Take as needed medications	%	64.1		62.4	66.9			78.6	71.0	58.7	60.3	65.0	66.9
	Medications for chronic conditions	%	30.5		31.1	27.6			33.8	28.4	32.3	29.7	31.0	30.9
	Medications to help quality of life	%	13.9		15.2	8.6			11.4	8.9	17.9	9.8	10.5	20.9
	Can't recall/unsure	%	1.2		.0	2.2			5.3	1.3	1.2	1.9	1.9	.0
	Refuse	%	7.1		5.4	6.9			.0	9.2	5.5	6.5	7.4	7.3
	Other	%	5.0		3.8	8.8			.0	2.7	6.7	5.6	1.9	7.1

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



# 2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

			Region							Gender		Age		
			Canada 2017-06	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 21 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] What was the one reason for [not filling/stopping early/taking a smaller dose of] your personal prescription on any occasion? [OPEN]	Total	Unwgt N	191	10	78	51	14	14	24	81	110	55	63	73
		Wgt N	194	9	79	53	14	16	24	91	103	66	61	67
	Too expensive/couldn't afford it	%	7.2		10.8	3.1				5.8	8.4	5.3	8.5	7.9
	Side effects	%	18.2		16.3	16.6				15.7	20.4	16.3	13.6	24.3
	No longer needed it	%	33.0		43.4	25.8				40.8	26.0	37.9	45.0	16.9
	No reason	%	.9		.0	1.6				.0	1.6	.0	.0	2.5
	I take it only when needed	%	5.2		3.9	6.3				4.4	5.8	1.8	4.3	9.3
	Meds were too strong	%	2.3		1.1	.0				3.0	1.7	4.1	.0	2.6
	Did not help my issues	%	2.5		2.3	5.6				3.4	1.7	.0	4.9	2.7
	Forgot/forgetful	%	7.0		5.7	8.7				5.5	8.4	12.7	3.1	5.1
	Allergic reaction	%	2.9		1.1	3.8				.0	5.5	3.5	1.4	3.7
	I prefer natural medicines	%	1.5		2.2	2.2				.0	2.8	1.8	2.8	.0
	Refuse	%	13.6		7.6	20.7				17.5	10.2	14.7	7.1	18.6
	I did not want to take it	%	2.0		1.1	1.6				1.4	2.4	.0	3.6	2.4
	Medication was changed	%	2.9		3.3	4.0				2.5	3.3	2.0	4.1	2.8
	Other	%	.5		1.1	.0				.0	.9	.0	.0	1.3
	Unsure	%	.5		.0	.0				.0	.9	.0	1.5	.0

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



## 2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

Question 22 - [IF COST IS THE REASON FOR PRESCRIPTION FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)? [OPEN]/ Question 23 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In the past six months, about how much was spent in total, at your own expense, on prescription medicines? [OPEN] \* Region

Mean		
	Question 22 - [IF COST IS THE REASON FOR PRESCRIPTION FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)? [OPEN]	Question 23 - [IF PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In the past six months, about how much was spent in total, at your own expense, on prescription medicines? [OPEN]
Total		270.35

\*Masked due to small sample size.

### Statistics

Question 23 - [IF PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In the past six months, about how much was spent in total, at your own expense, on prescription medicines?

N	Valid	147
	Missing	4297
Mean		270.35
Median		100.00

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 24 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] Are you aware or not aware of any publicly funded programs to help people access prescription medicines?	Total	Unwgt N	203	10*	83	55	15*	15*	25*	85	118	59	68	76
		Wgt N	206	9	84	57	15	16	24	96	110	71	66	69
	Aware	%	43.2		37.4	38.3				44.3	42.2	40.6	41.9	47.2
	Not aware	%	53.5		58.7	57.1				51.7	55.1	52.4	58.1	50.4
	Unsure	%	3.3		3.9	4.6				4.0	2.6	7.0	.0	2.5

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

			Region							Gender		Age		
			Canada											
			2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 25 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] I am going to read you a series of statements about your health benefit plan coverage and I would like you to tell me which best describes you? [RANDOMIZE]	Total	Unwgt N	203	10	83	55	15	15	25	85	118	59	68	76
		Wgt N	206	9	84	57	15	16	24	96	110	71	66	69
	I am covered through a work health benefit plan which is either my own or someone in my family	%	44.6		46.2	39.6				45.8	43.6	51.0	63.0	20.6
	I have health plan benefits and coverage which has been purchased, at our own expense through and insurance carrier	%	8.1		4.7	8.1				8.6	7.7	1.8	7.0	15.7
	I have health plan benefits and coverage through a union or professional association	%	12.8		12.4	20.6				15.9	10.1	18.8	14.3	5.3
	My prescription medicines are paid for by a government public plan but I am not a government employee	%	18.4		23.3	13.7				18.1	18.7	13.5	5.9	35.4
	I have no health plan coverage and all the time have to pay for the full price of any prescription I need	%	11.8		8.3	15.1				8.3	14.9	11.5	7.3	16.4
	Other	%	.8		.0	1.5				.0	1.5	.0	.0	2.4
	Refuse	%	3.4		5.0	1.4				3.3	3.4	3.4	2.5	4.2

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT

			Region							Gender		Age		
			Canada											
			2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Question 26 - [IF PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN AND NO INSURANCE FOR PRESCRIPTION MEDICINES] Why do you not have any insurance or government benefits for prescription medication? [OPEN]	Total	Unwgt N	27	0	7	10	2	3	5	7	20	7	5	15
		Wgt N	26	0	7	9	2	3	5	8	18	8	5	13
	Not available/offered to me	%												
	I don't need it	%												
	I am a senior	%												
	I am retired	%												
	I am not covered by my spouse's plan	%												
	I am unemployed	%												
	Don't want to pay more for insurance/Can't afford it	%												
	I have pension plan/other coverage	%												
	I am self-employed	%												
		Other	%											

\*Masked due to small sample size.

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – STAT SHEET - DRAFT**

			Region							Gender		Age		
			Canada 2017-07	Atlantic	Quebec	Ontario	Prairies	Alberta	BC	Male	Female	18 to 34	35 to 54	55 plus
Type	Total	Unwgt N	4445	312	1040	1720	302	491	580	1925	2520	1003	1582	1860
		Wgt N	4444	312	1041	1719	302	491	580	2148	2296	1212	1515	1717
	Short survey	%	75.0	80.7	73.8	75.6	76.8	77.3	69.5	71.9	77.9	63.1	78.4	80.5
	Long survey	%	4.8	3.0	6.9	3.9	5.1	4.5	4.4	4.5	5.0	6.9	4.4	3.6
	No prescription household	%	20.2	16.3	19.3	20.4	18.1	18.2	26.1	23.6	17.1	30.0	17.2	15.9

Nanos conducted an RDD dual frame (land- and cell- lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20. Subsamples with less than 30 responses have been masked.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 1 - In the past six months, has anyone in your household, including children, been given a prescription, either new or for a refill for a medication?	Total	Unwgt N	4445	3343	1102	204	454	471	443	397	409	431	176	188	1272
		Wgt N	4444	3429	1015	204	437	459	440	402	419	442	181	191	1270
	Yes	%	79.8	78.6	83.8	76.0	82.0	80.6	77.3	81.7	83.1	83.3	84.9	83.3	76.0
	No	%	20.2	21.4	16.2	24.0	18.0	19.4	22.7	18.3	16.9	16.7	15.1	16.7	24.0

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 2 - How many children under 18 years of age are in your household? [OPEN]	Total	Unwgt N	4445	3343	1102	204	454	471	443	397	409	431	176	188	1272
		Wgt N	4444	3429	1015	204	437	459	440	402	419	442	181	191	1270
	No/zero children	%	76.3	69.9	98.0	89.0	86.4	84.7	80.7	74.7	65.4	66.1	57.6	57.8	79.4
	One child	%	8.8	11.2	0.8	5.1	5.6	6.7	8.8	8.0	16.7	11.4	18.0	12.7	6.1
	Two children	%	9.2	12.0	0.0	3.4	4.7	4.5	5.9	11.7	11.2	13.4	18.2	20.1	8.9
	Three children	%	3.3	4.2	0.1	0.4	1.3	1.9	2.7	3.6	3.8	6.6	3.4	7.1	3.1
	Four children	%	1.2	1.5	0.0	0.0	1.1	1.0	0.8	1.3	2.2	1.8	2.8	1.1	0.7
	Refuse	%	1.2	1.2	1.2	2.2	1.0	1.3	1.1	0.8	0.7	0.7	0.0	1.2	1.7

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 1

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 3 - [IF ONE OR MORE CHILDREN] Have any of the children under 18 years of age been given a prescription in the past six months?	Total	Unwgt N	4445	3343	1102	204	454	471	443	397	409	431	176	188	1272
		Wgt N	4444	3429	1015	204	437	459	440	402	419	442	181	191	1270
	0 children	%	77.5	71.1	99.1	91.1	87.3	85.8	81.8	75.5	66.1	66.8	57.6	59.0	81.1
	Yes	%	13.3	17.1	0.3	4.6	8.0	8.8	9.5	15.3	16.0	22.6	26.8	23.8	11.1
	No	%	8.9	11.4	0.6	3.6	4.6	5.4	8.7	8.9	17.4	10.2	13.8	16.7	7.6
	Unsure	%	0.3	0.4	0.0	0.7	0.0	0.0	0.0	0.3	0.5	0.4	1.7	0.4	0.3

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 4 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS] All prescriptions for children under 18 years of age were taken without exception as prescribed?	Total	Unwgt N	555	552	3	9	31	38	38	58	66	93	45	45	132
		Wgt N	588	585			35	40	42	62	67	100	48	45	139
	Yes	%	94.2	94.3			94.2	94.5	94.5	94.3	97.0	95.5	94.1	89.0	93.0
	No	%	5.1	5.1			3.5	5.5	5.5	5.7	3.0	4.5	5.9	11.0	4.5
	Unsure	%	0.7	0.6			2.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.5

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 2

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65					Income							
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 5 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age received a prescription but the prescription was not filled (occasions in the past 6 months) [OPEN]	Total	Unwgt N	15	15	0	0	0	0	1	3	2	2	2	2	3
		Wgt N													
	1	%													
	2	%													
	4	%													
	Unsure	%													
	No answer	%													

\*Shaded due to small sample size.

			Over/Under 65					Income							
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 6 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age stopped take a prescriptions early (occasions in the past 6 months) [OPEN]	Total	Unwgt N	21	21	0	0	0	2	2	2	1	4	1	5	4
		Wgt N	23	23	0	0	0	2	2	2	1	4	1	5	4
	1	%													
	2	%													
	Unsure	%													

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 3

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65					Income							
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 7 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age took a smaller dose of the medication or less than prescribed (occasions in the past 6 months) [OPEN]	Total	Unwgt N	14	14	0	0	1	0	0	3	1	3	0	4	2
		Wgt N	15	15	0	0	1	0	0	3	1	3	0	4	2
	1	%													
	2	%													
	10	%													
	30	%													
	Unsure	%													

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 4

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 8 – [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND CHILD’S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In general, what were those prescriptions for? [OPEN]	Total	Unwgt N	22	22	0	0	1	1	2	3	2	3	2	3	5
		Wgt N	23	23	0	0	1	1	2	3	2	3	2	3	5
	Vitamin deficiency	%													
	Antibiotics	%													
	Asthma	%													
	ADHD	%													
	Acne medication	%													
	Migraines	%													
	Ear infection	%													
	Skin condition/ rash	%													
	Dyslexia	%													
	Mosquito bite	%													
	Chronic disease	%													
	Buccal infection	%													
	Lung infection	%													
	As a precaution	%													
	Anti-depressants	%													
	Tonsils	%													

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

		Over/Under 65				Income								
		Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Cold relief	%													
Allergies	%													
Refuse	%													

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 6

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 11 – [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPE D EARLY/ SMALLER DOSE TAKEN] What was the one reason for [not filling/stopping early/taking a smaller dose of] the prescription of the child under 18 years of age on any occasion? [OPEN]	Total	Unwgt N	28	28	0	0	1	2	2	3	2	4	3	5	6
		Wgt N	30	30	0	0	1	2	2	3	2	4	3	5	6
	Forgot/forgetful	%													
	We already had some	%													
	Wrong diagnosis	%													
	Their condition improved/no longer needed	%													
	To see how they would respond	%													
	Child chose not to take it	%													
	Doctor recommendation	%													
	It was not working	%													
	Allergic reaction to medication	%													
	It was unnecessary	%													
	Unsure	%													
	Refuse	%													

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 7

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

**Question 12 – [IF COST IS THE REASON A CHILD’S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN] \* Over/Under 65**

Mean	
Question 12 - [IF COST IS THE REASON A CHILD’S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN]	
Over/Under 65	
Under 65 years of age	
Total	

\*Shaded due to small sample size.

**Question 12 - [IF COST IS THE REASON A CHILD’S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN] \* Income**

Mean	
Question 12 - [IF COST IS THE REASON A CHILD’S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan? [OPEN]	
Income	
\$200,000 or more	
Total	

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 8

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 13 – [IF SOMEONE IN HOUSEHOLD HAS BEEN GIVEN PRESCRIPTION IN LAST SIX MONTHS] Have you been given a prescription in the past six months?	Total	Unwgt N	3590	2665	925	156	376	385	346	328	345	364	150	159	981
		Wgt N	3547	2696	851	155	358	370	340	329	349	368	153	159	965
	Yes	%	80.2	76.6	91.6	84.3	90.0	84.1	77.5	80.8	79.3	76.2	71.2	71.5	80.0
	No	%	19.6	23.3	8.1	15.7	10.0	15.9	22.5	19.2	20.7	23.8	28.8	28.5	19.5
	77	%	0.1	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3
	99	%	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 14 - [IF SOMEONE IN HOUSEHOLD HAS BEEN GIVEN PRESCRIPTION IN LAST SIX MONTHS AND PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS] You personally took all prescriptions without exception as prescribed	Total	Unwgt N	2904	2057	847	132	339	325	270	269	274	278	109	117	791
		Wgt N	2845	2065	780	130	323	311	264	265	276	281	109	113	772
	Yes	%	92.8	91.8	95.2	93.8	93.5	92.2	91.6	91.2	93.1	91.7	92.1	96.7	93.2
	No	%	7.2	8.2	4.8	6.2	6.5	7.8	8.4	8.8	6.9	8.3	7.9	3.3	6.8

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 9

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 15 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally stopped taking the prescription early (occasions in the past 6 months) [OPEN]	Total	Unwgt N	153	121	32	6	16	16	18	18	16	15	7	2	39
		Wgt N	157	127	29	6	16	15	18	18	17	16	8	2	40
	1	%	51.5	53.2	44.2										48.7
	2	%	28.9	27.8	33.8										25.7
	3	%	5.2	4.8	6.7										9.6
	4	%	0.8	1.0	0.0										3.2
	5	%	0.8	0.9	0.0										0.0
	6	%	0.8	1.0	0.0										3.2
	8	%	0.6	0.0	3.1										2.2
	10	%	1.5	1.0	3.3										0.0
	12	%	0.6	0.7	0.0										0.0
	15	%	0.8	1.0	0.0										0.0
	20	%	2.1	2.6	0.0										0.0
	30	%	0.8	1.0	0.0										0.0
	Unsure	%	5.7	4.9	8.9										7.2

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 10

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 16 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally received but did not fill a prescription (occasions in the past 6 months) [OPEN]	Total	Unwgt N	111	85	26	4	10	14	13	13	14	12	4	2	25
		Wgt N	112	88	24	4	10	14	13	14	15	12	4	2	24
	1	%	55.5	56.3											
	2	%	34.2	31.8											
	3	%	5.0	5.1											
	4	%	1.2	1.5											
	Unsure	%	1.9	2.5											
	Refuse	%	2.2	2.8											

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 11

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 17 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally took a smaller dose or less often than prescribed (occasions in the past 6 months) [OPEN]	Total	Unwgt N	145	114	31	5	16	21	17	15	12	15	4	4	36
		Wgt N	146	118	28	5	15	20	17	15	13	16	4	4	36
	1	%	43.4	44.3	39.7										32.4
	2	%	19.5	19.5	19.4										25.4
	3	%	2.0	0.0	10.5										0.0
	5	%	1.1	0.7	3.0										2.3
	6	%	1.2	0.7	3.5										0.0
	7	%	0.8	0.9	0.0										0.0
	10	%	6.4	7.9	0.0										3.0
	12	%	0.6	0.0	2.9										0.0
	15	%	0.8	0.9	0.0										0.0
	20	%	2.3	2.8	0.0										3.6
	21	%	0.6	0.0	3.0										0.0
	30	%	2.1	2.7	0.0										0.0
	Unsure	%	18.5	18.6	18.0										33.2
	Refuse	%	0.8	1.0	0.0										0.0

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 12

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 18 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In general, what were those personal prescriptions for? [OPEN]	Total	Unwgt N	178	141	37	8	20	19	20	21	16	22	8	3	41
		Wgt N	180	146	33	8	19	18	20	21	17	23	9	3	41
	Antibiotics	%	13.2	14.3	8.2										9.2
	Digestive issues	%	10.3	10.8	8.4										11.8
	Pain medication	%	20.6	21.6	16.3										21.3
	Infection	%	0.7	0.9	0.0										3.2
	Anxiety	%	5.3	5.9	2.8										5.9
	Incontinence	%	0.9	0.6	2.5										2.0
	Anti-inflammatory	%	9.1	9.5	7.7										11.7
	Depression	%	9.3	10.7	3.0										4.9
	Hypertension	%	3.3	2.2	8.4										2.2
	Allergies	%	4.9	6.1	0.0										2.0
	Fibromyalgia	%	0.5	0.6	0.0										0.0
	Can't recall	%	0.6	0.7	0.0										2.6
	ADHD	%	1.4	1.7	0.0										0.0
	Cholesterol	%	4.7	3.4	10.5										5.5
	Skin condition	%	2.6	2.5	3.0										0.0
	Osteoporosis	%	1.0	0.0	5.2										2.0
	Headache	%	0.5	0.6	0.0										2.1

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 13

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

		Over/Under 65			Income									Refuse/rather not say
		Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	
Athlete's Foot	%	0.5	0.0	2.9										0.0
Muscle spasms	%	2.6	3.2	0.0										0.0
Heart condition	%	0.5	0.0	2.5										0.0
Hemorrhoids	%	0.5	0.7	0.0										0.0
Rheumatoid arthritis	%	1.3	0.5	4.9										1.9
Abscessed tooth	%	0.5	0.6	0.0										0.0
Flu/cold	%	2.4	2.9	0.0										2.9
Eye drops	%	0.5	0.0	2.8										0.0
Puffer for asthma	%	2.2	1.5	5.2										4.2
Hormones	%	1.6	1.4	2.7										0.0
Sleep aids	%	4.6	5.7	0.0										0.0
Blood thinner	%	2.4	2.3	3.0										0.0
Alzheimer's	%	0.5	0.0	2.9										0.0
Blood pressure	%	4.4	4.2	5.2										4.3
Medication for lung condition	%	0.5	0.0	2.7										2.2
Medication for diabetes	%	4.5	4.9	2.7										3.2
Oral contraceptives	%	0.7	0.8	0.0										0.0
Hyperthyroidism	%	2.0	1.2	5.4										0.0

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 14

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

		Over/Under 65			Income									Refuse/rather not say
		Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	
Medication for after transplant surgery	%	1.0	1.2	0.0										2.1
Iron supplements	%	1.0	0.0	5.3										2.0
Central nervous system	%	0.5	0.0	2.7										0.0
Medicated shampoo	%	0.6	0.7	0.0										0.0
Fetal malformation	%	0.7	0.8	0.0										0.0
Other	%	5.6	5.0	7.9										9.0
Unsure	%	0.6	0.8	0.0										0.0
Refuse	%	7.6	7.6	7.9										17.2

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 15

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 21 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] What was the one reason for [not filling/stopping early/taking a smaller dose of] your personal prescription on any occasion? [OPEN]	Total	Unwgt N	191	150	41	8	22	21	20	22	18	22	8	3	47
		Wgt N	194	158	37	8	21	20	20	22	19	23	9	3	49
	Cost	%	7.2	6.6	9.6										8.4
	Side effects	%	18.2	17.8	19.7										18.6
	No longer needed it	%	33.0	37.7	12.6										23.0
	No reason	%	0.9	0.0	4.5										1.7
	I take it only when needed	%	5.2	4.7	7.2										10.6
	Meds were too strong	%	2.3	2.2	2.4										2.6
	Did not help my issues	%	2.5	2.5	2.3										3.5
	Forgot/forgetful	%	7.0	7.6	4.7										7.8
	Allergic reaction	%	2.9	2.5	4.5										5.2
	I prefer natural medicines	%	1.5	1.8	0.0										0.0
	Refuse	%	13.6	10.8	25.6										16.9
	I did not want to take it	%	2.0	1.4	4.4										1.6
	Medication was changed	%	2.9	3.0	2.4										0.0
	Other	%	0.5	0.6	0.0										0.0
	Unsure	%	0.5	0.6	0.0										0.0

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 16

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

**Question 22 - [IF COST IS THE REASON FOR PRESCRIPTION FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)? [OPEN]/ Question 23 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In the past six months, about how much was spent in total, at your own expense, on prescription medicines? [OPEN] \* Over/Under 65**

Mean		
	Question 22 - [IF COST IS THE REASON FOR PRESCRIPTION FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)? [OPEN]	Question 23 - [IF PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In the past six months, about how much was spent in total, at your own expense, on prescription medicines? [OPEN]
Over/Under 65		
Under 65 years of age		281.86
65 and older		224.49
Total		270.35

\*Shaded due to small sample size.

**Question 22 - [IF COST IS THE REASON FOR PRESCRIPTION FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)? [OPEN]/ Question 23 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In the past six months, about how much was spent in total, at your own expense, on prescription medicines? [OPEN] \* Income**

Mean		
	Question 22 - [IF COST IS THE REASON FOR PRESCRIPTION FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)? [OPEN]	Question 23 - [IF PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In the past six months, about how much was spent in total, at your own expense, on prescription medicines? [OPEN]
Income		
\$0 to \$14,999		57.12
\$15,000 to \$29,999		555.63
\$30,000 to \$44,999		118.68
\$45,000 to \$59,999		277.53
\$60,000 to \$74,999		253.37
\$75,000 to \$99,999		379.29
\$100,000 to \$149,999		110.92
\$150,000 to \$199,999		843.31
Refuse/rather not say		144.55
Total		270.35

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 17

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 24 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] Are you aware or not aware of any publicly funded programs to help people access prescription medicines?	Total	Unwgt N	203	161	42	8	22	25	22	23	18	22	8	4	51
		Wgt N	206	169	38	8	21	24	22	23	19	23	9	4	53
	Aware	%	43.2	42.7	45.5										39.7
	Not aware	%	53.5	53.8	52.1										53.0
	Unsure	%	3.3	3.5	2.4										7.3

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 18

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 25 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] I am going to read you a series of statements about your health benefit plan coverage and I would like you to tell me which best describes you? [RANDOMIZE]	Total	Unwgt N	203	161	42	8	22	25	22	23	18	22	8	4	51
		Wgt N	206	169	38	8	21	24	22	23	19	23	9	4	53
	I am covered through a work health benefit plan which is either my own or someone in my family	%	44.6	52.6	8.7										44.0
	I have health plan benefits and coverage which has been purchased, at our own expense through and insurance carrier	%	8.1	5.5	19.7										9.2
	I have health plan benefits and coverage through a union or professional association	%	12.8	14.0	7.6										11.8
	My prescription medicines are paid for by a government public plan but I am not a government employee	%	18.4	12.0	47.3										20.2
	I have no health plan coverage and all the time have to pay for the full price of any prescription I need	%	11.8	11.9	11.6										8.5
	Other	%	0.8	1.0	0.0										0.0
	Refuse	%	3.4	3.0	5.1										6.2

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 19

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – AGE + INCOME - INTERNAL USE ONLY - STAT SHEET - DRAFT**

			Over/Under 65			Income									
			Canada 2017-06	Under 65 years of age	65 and older	\$0 to \$14,999	\$15,000 to \$29,999	\$30,000 to \$44,999	\$45,000 to \$59,999	\$60,000 to \$74,999	\$75,000 to \$99,999	\$100,000 to \$149,999	\$150,000 to \$199,999	\$200,000 or more	Refuse/rather not say
Question 26 - [IF PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN AND NO INSURANCE FOR PRESCRIPTION MEDICINES] Why do you not have any insurance or government benefits for prescription medication? [OPEN]	Total	Unwgt N	27	22	5	2	7	3	4	4	1	0	0	1	5
		Wgt N	26	22	4	2	6	3	4	4	1	0	0	1	4
	Not available/offered to me	%													
	I don't need it	%													
	I am a senior	%													
	I am retired	%													
	I am not covered by my spouse's plan	%													
	I am unemployed	%													
	Don't want to pay more for insurance/Can't afford it	%													
	I have pension plan/other coverage	%													
	I am self-employed	%													
	Other	%													

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 20

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income				
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+
Type	Total	Unwgt N	4445	658	914	806	795
		Wgt N	4444	640	899	821	813
	Short survey	%	75.0	76.2	74.2	77.1	78.0
	Long survey	%	4.8	3.9	4.8	5.3	5.7
	No prescription household	%	20.2	19.9	21.0	17.5	16.3

			Grouped Income				
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+
Question 1 - In the past six months, has anyone in your household, including children, been given a prescription, either new or for a refill for a medication?	Total	Unwgt N	4445	658	914	806	795
		Wgt N	4444	640	899	821	813
	Yes	%	79.8	80.1	79.0	82.5	83.7
	No	%	20.2	19.9	21.0	17.5	16.3

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 1

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 2 - How many children under 18 years of age are in your household? [Open-ended]	Total	Unwgt N	4445	658	914	806	795	1272
		Wgt N	4444	640	899	821	813	1270
	No/zero children	%	76.3	87.2	82.8	69.9	62.3	79.4
	One child	%	8.8	5.4	7.7	12.4	13.2	6.1
	Two children	%	9.2	4.3	5.2	11.4	16.0	8.9
	Three children	%	3.3	1.0	2.2	3.7	6.0	3.1
	Four children	%	1.2	.7	.9	1.8	1.9	.7
	Refuse	%	1.2	1.3	1.2	.7	.7	1.7

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 3 - [IF ONE OR MORE CHILDREN] Have any of the children under 18 years of age been given a prescription in the past six months?	Total	Unwgt N	4445	658	914	806	795	1272
		Wgt N	4444	640	899	821	813	1270
	0 children	%	77.5	88.5	83.9	70.7	62.9	81.1
	Yes	%	13.3	6.9	9.1	15.7	23.8	11.1
	No	%	8.9	4.3	7.0	13.3	12.5	7.6
	Unsure	%	.3	.2	.0	.4	.7	.3

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 2

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 4 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS] All prescriptions for children under 18 years of age were taken without exception as prescribed?	Total	Unwgt N	555	40	76	124	183	132
		Wgt N	588	44	82	129	194	139
	Yes	%	94.2	95.4	94.5	95.7	93.6	93.0
	No	%	5.1	2.7	5.5	4.3	6.4	4.5
	Unsure	%	.7	1.9	.0	.0	.0	2.5

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 5 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age received a prescription but the prescription was not filled (occasions in the past 6 months) [OPEN]	Total	Unwgt N	15	0	1	5	6	3
		Wgt N	17	0	1	6	6	4
	1	%						
	2	%						
	4	%						
	Unsure	%						
	No answer	%						

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 3

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 6 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age stopped take a prescriptions early (occasions in the past 6 months) [OPEN]	Total	Unwgt N	21	0	4	3	10	4
		Wgt N	23	0	5	4	11	4
	1	%						
	2	%						
	Unsure	%						

\*Shaded due to small sample size.

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 7 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION WAS NOT TAKEN AS PRESCRIBED] A child under 18 years of age took a smaller dose of the medication or less than prescribed (occasions in the past 6 months) [OPEN]	Total	Unwgt N	14	1	0	4	7	2
		Wgt N	15	1	0	5	7	2
	1	%						
	2	%						
	10	%						
	30	%						
	Unsure	%						

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 4

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 8 – [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In general, what were those prescriptions for? [OPEN]	Total	Unwgt N	22	1	3	5	8	5
		Wgt N	23	1	3	6	8	5
	Vitamin deficiency	%						
	Antibiotics	%						
	Asthma	%						
	ADHD	%						
	Acne medication	%						
	Migraines	%						
	Ear infection	%						
	Skin condition/rash	%						
	Dyslexia	%						
	Mosquito bite	%						
	Chronic disease	%						
	Buccal infection	%						
	Lung infection	%						
	As a precaution	%						
	Anti-depressants	%						
	Tonsils	%						
	Cold relief	%						

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 5

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

		Grouped Income					
		Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Allergies	%						
Refuse	%						

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 6

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 11 - [IF CHILD RECEIVED PRESCRIPTION IN PAST SIX MONTHS AND CHILD'S PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] What was the one reason for [not filling/stopping early/taking a smaller dose of] the prescription of the child under 18 years of age on any occasion? [OPEN]	Total	Unwgt N	28	1	4	5	12	6
		Wgt N	30	1	5	6	12	6
	Forgot/forgetful	%						
	We already had some	%						
	Wrong diagnosis	%						
	Their condition improved/no longer needed	%						
	To see how they would respond	%						
	Child chose not to take it	%						
	Doctor recommendation	%						
	It was not working	%						
	Allergic reaction to medication	%						
	It was unnecessary	%						
	Unsure	%						
	Refuse	%						

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 7

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

**Question 12 - [IF COST IS THE REASON A CHILD'S  
PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/  
SMALLER DOSE TAKEN] How much did that prescription  
cost to you or your family excluding any portion of the cost  
paid by a private drug plan? [OPEN] \* \$100,00 +**

Mean	
	Question 12 - [If cost] How much did that prescription cost to you or your family excluding any portion of the cost paid by a private drug plan?
Grouped Income	
\$100,000+	
Total	

\*Shaded due to small sample size.

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 13 - [IF SOMEONE IN HOUSEHOLD HAS BEEN GIVEN PRESCRIPTION IN LAST SIX MONTHS] Have you been given a prescription in the past six months	Total	Unwgt N	3590	532	731	673	673	981
		Wgt N	3547	513	710	677	680	965
	Yes	%	80.2	88.3	80.9	80.0	74.0	80.0
	No	%	19.6	11.7	19.1	20.0	26.0	19.5
	77	%	.1	.0	.0	.0	.0	.3
	99	%	.1	.0	.0	.0	.0	.3

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 8

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 14 - [IF SOMEONE IN HOUSEHOLD HAS BEEN GIVEN PRESCRIPTION IN LAST SIX MONTHS AND PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS] You personally took all prescriptions without exception as prescribed	Total	Unwgt N	2904	471	595	543	504	791
		Wgt N	2845	453	575	542	503	772
	Yes	%	92.8	93.6	91.9	92.2	92.9	93.2
	No	%	7.2	6.4	8.1	7.8	7.1	6.8

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 9

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 15 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally stopped taking the prescription early (occasions in the past 6 months) [OPEN]	Total	Unwgt N	153	22	34	34	24	39
		Wgt N	157	22	33	35	26	40
	1	%	51.5		41.7	52.3		48.7
	2	%	28.9		44.1	22.9		25.7
	3	%	5.2		3.0	5.9		9.6
	4	%	.8		.0	.0		3.2
	5	%	.8		.0	3.3		.0
	6	%	.8		.0	.0		3.2
	8	%	.6		.0	.0		2.2
	10	%	1.5		.0	3.7		.0
	12	%	.6		2.8	.0		.0
	15	%	.8		.0	.0		.0
	20	%	2.1		.0	9.3		.0
	30	%	.8		.0	.0		.0
	Unsure	%	5.7		8.4	2.5		7.2

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 10

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 16 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally received but did not fill a prescription (occasions in the past 6 months) [OPEN]	Total	Unwgt N	111	14	27	27	18	25
		Wgt N	112	14	26	29	18	24
	1	%	55.5					
	2	%	34.2					
	3	%	5.0					
	4	%	1.2					
	Unsure	%	1.9					
	Refuse	%	2.2					

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 11

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 17 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] You personally took a smaller dose or less often than prescribed (occasions in the past 6 months) [OPEN]	Total	Unwgt N	145	21	38	27	23	36
		Wgt N	146	20	38	28	24	36
	1	%	43.4		45.4			32.4
	2	%	19.5		24.1			25.4
	3	%	2.0		2.4			.0
	5	%	1.1		.0			2.3
	6	%	1.2		.0			.0
	7	%	.8		.0			.0
	10	%	6.4		5.3			3.0
	12	%	.6		.0			.0
	15	%	.8		.0			.0
	20	%	2.3		.0			3.6
	21	%	.6		.0			.0
	30	%	2.1		4.8			.0
	Unsure	%	18.5		18.1			33.2
	Refuse	%	.8		.0			.0

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 12

Internal Use Only



2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Question 18 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In general, what were those personal prescriptions for? [OPEN]	Total	Unwgt N	178	28	39	37	33	41
		Wgt N	180	27	38	38	35	41
	Antibiotics	%	13.2		14.7	11.3	22.2	9.2
	Digestive issues	%	10.3		9.1	3.1	15.6	11.8
	Pain medication	%	20.6		28.0	22.5	11.0	21.3
	Infection	%	.7		.0	.0	.0	3.2
	Anxiety	%	5.3		3.1	2.4	2.7	5.9
	Incontinence	%	.9		.0	.0	.0	2.0
	Anti-inflammatory	%	9.1		7.5	14.8	8.9	11.7
	Depression	%	9.3		12.6	5.2	3.2	4.9
	Hypertension	%	3.3		5.2	5.7	.0	2.2
	Allergies	%	4.9		.0	15.1	3.2	2.0
	Fibromyalgia	%	.5		2.3	.0	.0	.0
	Can't recall	%	.6		.0	.0	.0	2.6
	ADHD	%	1.4		.0	.0	3.8	.0
	Cholesterol	%	4.7		2.3	5.2	4.8	5.5
	Skin condition	%	2.6		2.3	2.6	5.3	.0
	Osteoporosis	%	1.0		.0	.0	.0	2.0
	Refuse	%	7.0		8.6	2.3	3.8	14.5

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 13

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Headache	%		.5		.0	.0	.0	2.1
Athlete's Foot	%		.5		.0	2.5	.0	.0
Muscle spasms	%		2.6		2.6	.0	10.7	.0
Heart condition	%		.5		.0	.0	.0	.0
Hemorrhoids	%		.5		.0	.0	2.8	.0
Rheumatoid arthritis	%		1.3		2.0	.0	.0	1.9
Abscessed tooth	%		.5		2.4	.0	.0	.0
Flu/cold	%		2.4		2.1	.0	6.5	2.9
Eye drops	%		.5		2.4	.0	.0	.0
Puffer for asthma	%		2.2		2.4	3.4	.0	4.2
Hormones	%		1.6		4.6	3.1	.0	.0
Sleep aids	%		4.6		.0	14.0	6.1	.0
Blood thinner	%		2.4		2.1	.0	7.2	.0
Alzheimer's	%		.5		.0	2.5	.0	.0
Blood pressure	%		4.4		3.2	2.1	3.5	4.3
Medication for lung condition	%		.5		.0	.0	.0	2.2
Medication for diabetes	%		4.5		2.9	5.2	5.7	3.2
Oral contraceptives	%		.7		.0	3.1	.0	.0
Hyperthyroidism	%		2.0		2.5	.0	.0	.0

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 14

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

		Grouped Income					
		Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+	Refuse
Medication for after transplant surgery	%	1.0		2.3	.0	.0	2.1
Iron supplements	%	1.0		.0	.0	.0	2.0
Central nervous system	%	.5		2.3	.0	.0	.0
Medicated shampoo	%	.6		.0	.0	2.9	.0
Fetal malformation	%	.7		.0	3.1	.0	.0
Other	%	5.6		2.3	2.2	6.9	9.0
Unsure	%	.6		.0	3.0	.0	.0
Refuse	%	.6		.0	.0	.0	2.7

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 15

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,000 to \$99,999	\$100,000+	Refuse
Question 21 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] What was the one reason for [not filling/stopping early/taking a smaller dose of] your personal prescription on any occasion? [OPEN]	Total	Unwgt N	191	30	41	40	33	47
		Wgt N	194	29	40	42	35	49
	Cost	%	7.2		7.2	10.2	.0	8.4
	Side effects	%	18.2		20.7	9.3	20.4	18.6
	No longer needed it	%	33.0		29.2	39.4	51.1	23.0
	No reason	%	.9		.0	.0	.0	1.7
	I take it only when needed	%	5.2		.0	4.4	2.8	10.6
	Meds were too strong	%	2.3		5.7	2.0	.0	2.6
	Did not help my issues	%	2.5		.0	2.4	3.0	3.5
	Forgot/forgetful	%	7.0		10.2	2.8	5.9	7.8
	Allergic reaction	%	2.9		2.9	2.8	2.2	5.2
	I prefer natural medicines	%	1.5		2.1	.0	5.9	.0
	Refuse	%	13.6		17.5	10.0	6.2	16.9
	I did not want to take it	%	2.0		.0	7.3	.0	1.6
	Medication was changed	%	2.9		4.5	5.1	2.5	.0
	Other	%	.5		.0	2.1	.0	.0
	Unsure	%	.5		.0	2.2	.0	.0

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is ±1.4 percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 16

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

**Question 22 - [IF COST IS THE REASON FOR PRESCRIPTION FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)? [OPEN]/ Question 23 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] In the past six months, about how much was spent in total, at your own expense, on prescription medicines? [OPEN]**

Mean		
Grouped Income	Question 22 - [If cost] - How much did that prescription cost to you or your family (excluding any portion of the cost paid by a private drug plan)?	Question 23 - In the past six months, about how much was spent in total, at your own expense, on prescription medicines?
\$0 to \$29,999		
\$30,000 to \$59,999		205.19
\$60,00 to \$99,999		306.64
\$100,000+		330.96
Refuse		144.55
Total		270.35

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 17

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,000 to \$99,999	\$100,000+	Refuse
Question 24 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] Are you aware or not aware of any publicly funded programs to help people access prescription medicines?	Total	Unwgt N	203	30	47	41	34	51
		Wgt N	206	29	46	43	36	53
	Aware	%	43.2		32.5	56.6	46.9	39.7
	Not aware	%	53.5		67.5	43.4	53.1	53.0
	Unsure	%	3.3		.0	.0	.0	7.3

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 18

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income					
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,000 to \$99,999	\$100,000+	Refuse
Question 25 - [IF PERSONALLY RECEIVED A PRESCRIPTION IN PAST SIX MONTHS AND PRESCRIPTION NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN] I am going to read you a series of statements about your health benefit plan coverage and I would like you to tell me which best describes you? [RANDOMIZE]	Total	Unwgt N	203	30	47	41	34	51
		Wgt N	206	29	46	43	36	53
	I am covered through a work health benefit plan which is either my own or someone in my family	%	44.6		37.6	60.9	63.4	44.0
	I have health plan benefits and coverage which has been purchased, at our own expense through and insurance carrier	%	8.1		5.8	12.8	7.8	9.2
	I have health plan benefits and coverage through a union or professional association	%	12.8		19.1	8.3	15.5	11.8
	My prescription medicines are paid for by a government public plan but I am not a government employee	%	18.4		18.9	5.2	5.4	20.2
	I have no health plan coverage and all the time have to pay for the full price of any prescription I need	%	11.8		14.8	12.8	.0	8.5
	Other	%	.8		.0	.0	2.4	.0
	Refuse	%	3.4		3.7	.0	5.5	6.2

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 19

Internal Use Only



**2017-1025 – IMC and CLHIA – Prescription Use Survey – Grouped Income Crosstab - STAT SHEET - DRAFT**

			Grouped Income				
			Canada 2017-06	\$0 to \$29,999	\$30,000 to \$59,999	\$60,00 to \$99,999	\$100,000+
Question 26 - [IF PRESCRIPTION WAS NOT FILLED/STOPPED EARLY/ SMALLER DOSE TAKEN AND NO INSURANCE FOR PRESCRIPTION MEDICINES] Why do you not have any insurance or government benefits for prescription medication? [OPEN]	Total	Unwgt N	27	9	7	5	1
		Wgt N	26	8	7	5	1
	Not available/offered to me	%					
	I don't need it	%					
	I am a senior	%					
	I am retired	%					
	I am not covered by my spouse's plan	%					
	I am unemployed	%					
	Don't want to pay more for insurance/Can't afford it	%					
	I have pension plan/other coverage	%					
	I am self-employed	%					
Other		%					

\*Shaded due to small sample size.

Nanos conducted an RDD dual frame (land- and cell-lines) random telephone survey of 4445 Canadians, 18 years of age or older, between May 27<sup>th</sup> to July 10<sup>th</sup>, 2017. The margin of error for a random survey of 4445 Canadians is  $\pm 1.4$  percentage points, 19 times out of 20.

[www.nanosresearch.com](http://www.nanosresearch.com)

Page 20

Internal Use Only